

**Overdose Response:
SAVE ME**

- ❖ Stimulate: Speak loudly, rubbing sternum & pinching skin. Unresponsive? Call 911.
- ❖ Airway. Check airway: tilt head and lift chin to open.
- ❖ Ventilate: 1 breath every 5 seconds, using CPR mask. If their heart has stopped, administer CPR to tune of “Staying Alive” (100 compressions/minute)
- ❖ Evaluate: Are they now breathing?

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME (cont.)**

- ❖ Medication: 1 dose naloxone (spray in nostrils or direct IM dosage) & write down time.
- ❖ Evaluate & support. Naloxone takes 2-5 minutes to work. Wait 3-5 minutes while continuing rescue breaths (approximately 40). If no change, administer second naloxone dose.

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME**

- ❖ Stimulate: Speak loudly, rubbing sternum & pinching skin. Unresponsive? Call 911.
- ❖ Airway. Check airway: tilt head and lift chin to open.
- ❖ Ventilate: 1 breath every 5 seconds, using CPR mask. If their heart has stopped, administer CPR to tune of “Staying Alive” (100 compressions/minute)
- ❖ Evaluate: Are they now breathing?

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME (cont.)**

- ❖ Medication: 1 dose naloxone (spray in nostrils or direct IM dosage) & write down time.
- ❖ Evaluate & support. Naloxone takes 2-5 minutes to work. Wait 3-5 minutes while continuing rescue breaths (approximately 40). If no change, administer second naloxone dose.

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME**

- ❖ Stimulate: Speak loudly, rubbing sternum & pinching skin. Unresponsive? Call 911.
- ❖ Airway. Check airway: tilt head and lift chin to open.
- ❖ Ventilate: 1 breath every 5 seconds, using CPR mask. If their heart has stopped, administer CPR to tune of “Staying Alive” (100 compressions/minute)
- ❖ Evaluate: Are they now breathing?

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME (cont.)**

- ❖ Medication: 1 dose naloxone (spray in nostrils or direct IM dosage) & write down time.
- ❖ Evaluate & support. Naloxone takes 2-5 minutes to work. Wait 3-5 minutes while continuing rescue breaths (approximately 40). If no change, administer second naloxone dose.

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME**

- ❖ Stimulate: Speak loudly, rubbing sternum & pinching skin. Unresponsive? Call 911.
- ❖ Airway. Check airway: tilt head and lift chin to open.
- ❖ Ventilate: 1 breath every 5 seconds, using CPR mask. If their heart has stopped, administer CPR to tune of “Staying Alive” (100 compressions/minute)
- ❖ Evaluate: Are they now breathing?

© 2022 Psychological Safety Alliance

**Overdose Response:
SAVE ME (cont.)**

- ❖ Medication: 1 dose naloxone (spray in nostrils or direct IM dosage) & write down time.
- ❖ Evaluate & support. Naloxone takes 2-5 minutes to work. Wait 3-5 minutes while continuing rescue breaths (approximately 40). If no change, administer second naloxone dose.

© 2022 Psychological Safety Alliance

How to Recognize an Opioid Overdose


- Not moving & not responsive to attempts to awaken
- Slow, erratic, or no breathing (under 10 breaths/minute)
- Slow, erratic, or no hearttrate (under 50 beats/minute)
- Tiny, "pinpoint" pupils
- Blue/gray lips & nailbeds
- Cold, clammy skin

© 2023 PSYCHEDELIC SAFETY ALLIANCE

How to Recognize an Opioid Overdose


- Not moving & not responsive to attempts to awaken
- Slow, erratic, or no breathing (under 10 breaths/minute)
- Slow, erratic, or no hearttrate (under 50 beats/minute)
- Tiny, "pinpoint" pupils
- Blue/gray lips & nailbeds
- Cold, clammy skin

© 2023 PSYCHEDELIC SAFETY ALLIANCE



If you have to leave the person unattended to seek help, place in recovery position. Naloxone only temporarily reverses an overdose, and wears off within 20-90 minutes. They can still overdose. Stay with the person until EMTs arrive. Request help from other staff or bystanders if needed.

© 2023 PSYCHEDELIC SAFETY ALLIANCE



If you have to leave the person unattended to seek help, place in recovery position. Naloxone only temporarily reverses an overdose, and wears off within 20-90 minutes. They can still overdose. Stay with the person until EMTs arrive. Request help from other staff or bystanders if needed.

© 2023 PSYCHEDELIC SAFETY ALLIANCE

How to Recognize an Opioid Overdose


- Not moving & not responsive to attempts to awaken
- Slow, erratic, or no breathing (under 10 breaths/minute)
- Slow, erratic, or no hearttrate (under 50 beats/minute)
- Tiny, "pinpoint" pupils
- Blue/gray lips & nailbeds
- Cold, clammy skin

© 2023 PSYCHEDELIC SAFETY ALLIANCE

How to Recognize an Opioid Overdose


- Not moving & not responsive to attempts to awaken
- Slow, erratic, or no breathing (under 10 breaths/minute)
- Slow, erratic, or no hearttrate (under 50 beats/minute)
- Tiny, "pinpoint" pupils
- Blue/gray lips & nailbeds
- Cold, clammy skin

© 2023 PSYCHEDELIC SAFETY ALLIANCE



If you have to leave the person unattended to seek help, place in recovery position. Naloxone only temporarily reverses an overdose, and wears off within 20-90 minutes. They can still overdose. Stay with the person until EMTs arrive. Request help from other staff or bystanders if needed.

© 2023 PSYCHEDELIC SAFETY ALLIANCE



If you have to leave the person unattended to seek help, place in recovery position. Naloxone only temporarily reverses an overdose, and wears off within 20-90 minutes. They can still overdose. Stay with the person until EMTs arrive. Request help from other staff or bystanders if needed.

© 2023 PSYCHEDELIC SAFETY ALLIANCE